



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

148. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College Defiance, Ohio.

Helmholtz's differential equation for the strength of an electric current C at any time t , is $C=E/R-L/R \times dC/dt$. Solve this equation, supposing $C=0$ when $t=0$; and E, R, L are to be regarded as constants.

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa., and J. SCHEFFER, A. M. Hagerstown, Md.

The equation can be written $LdC=(E-CR)dt$.

$$\therefore \frac{LdC}{E-CR}=dt, \text{ or } \frac{L}{R} \log(E-CR)+t+B=0.$$

$$\text{When } C=0, t=0, \text{ and } B=-\frac{L}{R} \log E.$$

$$\therefore t + \frac{L}{R} \log(E-CR) = \frac{L}{R} \log E, \text{ or } t + \frac{L}{R} \log \left(\frac{E-CR}{E} \right) = 0.$$

149. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Find the volume contained between the plane $z=(a-x)\cot\beta$ and the surface $xz^2=(a-x)(x^2-y^2)$.

Solution by the PROPOSER.

$$\text{The } z \text{ limits are } z=\sqrt{\frac{a-x}{x} (x^2+y^2)} \text{ to } z=(a-x)\cot\beta.$$

$$\text{Eliminating } z, y=\pm\sqrt{[x(a\cot^2\beta-x\operatorname{cosec}^2\beta)]}=y'.$$

$$\text{The } x \text{ limits are } 0 \text{ and } a\cos^2\beta=x'.$$

$$\begin{aligned} V &= 2 \int_0^{x'} \int_0^{y'} [(a-x)\cot\beta - \sqrt{\frac{a-x}{x} (x^2+y^2)}] dx dy \\ &= \int_0^{x'} \left[\sqrt{x} (a-x)\cot\beta \sqrt{a\cot^2\beta - x\operatorname{cosec}^2\beta} \right. \\ &\quad \left. - x \sqrt{(a-x)^2} \log \left(\frac{\sqrt{(a-x)\cot\beta} + \sqrt{(a\cot^2\beta - x\operatorname{cosec}^2\beta)}}{\sqrt{x}} \right) \right] dx. \end{aligned}$$

$$\text{Let } x=a\cos^2\beta\sin^2\theta.$$

$$\therefore V=2a^3\cos^3\beta\cot^2\beta \int_0^{\frac{1}{2}\pi} (1-\cos^2\beta\sin^2\theta) \sin^2\theta \cos^2\theta d\theta$$

$$-2a^3\cos^5\beta \int_0^{\frac{1}{2}\pi} \sqrt{1-\cos^2\beta\sin^2\theta} \sin^4\theta \cos\theta \log \left(\frac{\sqrt{1-\cos^2\beta\sin^2\theta} + \cos\theta}{\sin\beta\sin\theta} \right) d\theta.$$

Let $\cos\beta\sin\theta = \sin\varphi$; then the second term becomes

$$2a^3 \int_0^{\frac{1}{2}\pi - \beta} \sin^4 \varphi \cos^2 \varphi \log \left[\frac{\cos \varphi \cos \beta + \sqrt{(\cos^2 \beta - \sin^2 \varphi)}}{\sin \beta \sin \varphi} \right] d\varphi = 2a^3 \int_0^{\frac{1}{2}\pi - \beta} \left(\frac{1}{16} \varphi - \frac{1}{16} \sin \varphi \cos \varphi - \frac{1}{4} \sin^3 \varphi \cos \varphi + \frac{1}{6} \sin^5 \varphi \cos \varphi \right) \frac{\cos \beta d\varphi}{\sin \varphi \sqrt{(\cos^2 \beta - \sin^2 \varphi)}}.$$

$$\therefore V = \frac{1}{16} \pi a^3 \cos^3 \beta \cot^2 \beta (1 + \sin^2 \beta) + \frac{1}{8} a^3 \cos \beta \int_0^{\frac{1}{2}\pi} d\theta + \frac{1}{2} \cos^3 \beta \int_0^{\frac{1}{2}\pi} \sin^2 \theta d\theta - \frac{1}{8} a^3 \cos^5 \beta \int_0^{\frac{1}{2}\pi} \sin^4 \theta d\theta - \frac{1}{8} a^3 \int_0^{\frac{1}{2}\pi} \frac{\sin^{-1}(\cos \beta \sin \theta) d\theta}{\sin \theta \sqrt{(1 - \cos^2 \beta \sin^2 \theta)}}.$$

$$\therefore V = \frac{1}{16} \pi a^3 \cot \beta \operatorname{cosec} \beta - \frac{1}{4} \pi a^3 \cos^3 \beta$$

$$- \frac{1}{8} a^3 \int_0^{\frac{1}{2}\pi} \frac{\sin^{-1}(\cos \beta \sin \theta) d\theta}{\sin \theta \sqrt{(1 - \cos^2 \beta \sin^2 \theta)}} - \frac{1}{8} a^3 \int_0^{\frac{1}{2}\pi} \frac{\sin^{-1}(\cos \beta \sin \theta) d\theta}{\sin \theta \sqrt{(1 - \cos^2 \beta \sin^2 \theta)}}$$

$$= \frac{1}{8} a^3 \cos \beta \int_0^{\frac{1}{2}\pi} (1 + \frac{2}{3} \cos^2 \beta \sin^2 \theta + \frac{8}{15} \cos^4 \beta \sin^4 \theta + \frac{16}{35} \cos^6 \beta \sin^6 \theta + \dots) d\theta$$

$$= \frac{1}{16} \pi a^3 (\cos \beta + \frac{1}{3} \cos^3 \beta + \frac{1}{5} \cos^5 \beta + \frac{1}{7} \cos^7 \beta + \dots) = \frac{1}{16} \pi a^3 \log \cot \frac{1}{2} \beta.$$

$$\therefore V = \frac{1}{4} \pi a^3 (3 \cot \beta \operatorname{cosec} \beta - 2 \cos^3 \beta - 3 \log \cot \frac{1}{2} \beta).$$


150. Proposed by E. B. ESCOTT, Instructor in Mathematics, University of Michigan, Ann Arbor, Mich.

Find total area between the curve $x^4 y - x^2 + 4y - 1 = 0$ and the x -axis.

Solution by J. E. SANDERS, Hackney, Ohio, and the PROPOSER.

The equation may be written $y = \frac{x^2 + 1}{x^4 + 4}$.

$$\text{Area} = 2 \int_0^\infty y dx = 2 \int_0^\infty \frac{x^2 + 1}{x^4 + 4} dx$$



$$= \frac{1}{4} \left[\int_0^\infty \frac{x+2}{x^2-2x+2} dx - \int_0^\infty \frac{x-2}{x^2+2x+2} dx \right] = \frac{1}{8} \left[\log(x^2-2x+2) \right.$$

$$\left. - \log(x^2+2x+2) + 6[\tan^{-1}(x+1) + \tan^{-1}(x-1)] \right]_0^\infty$$

$$= \frac{1}{8} \left[\log \frac{x^2-2x+2}{x^2+2x+2} + 6 \tan^{-1} \frac{2x}{2-x^2} \right]_0^\infty = \frac{1}{8} (6\pi) = \frac{3}{4} \pi. \quad \text{Answer.}$$